

Case Report

Compound Odontoma in the Anterior Mandible: Diagnostic Challenges and Surgical Management of a Rare Presentation

Vikram Karande¹. Sanjay Ranade². Vaishali Pagare¹

¹ Department of Oral & Maxillofacial Surgery, D.Y. Patil Dental School, Lohegaon, Pune, Maharashtra, India

² Department of General Surgery, D.Y. Patil Dental School, Lohegaon, Pune, Maharashtra, India

Citation: Karande V. Odontoma in the Anterior Mandible: Diagnostic Challenges and Surgical Management of a Rare Presentation. Oral Sphere J Dent Health Sci. (2025); 1(3): 173-177

For reprints contact:

publisher@fontfusionspublication.com

Received: March 12, 2025;

Revised: April 15, 2025;

Accepted: May 08, 2025;

Published: July 01, 2025

***Corresponding author:** Vikram Karande
Email: drvikramkarande@gmail.com

DOI: <https://doi.org/10.63150/osidhs.2025.15>

© The Author(s). (2025)

Published by Font Fusions Publication

Open Access.

This article is licensed under the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), allowing non-commercial sharing, adaptation, and distribution, provided you give appropriate credit to the original author(s), provide a link to the license, and indicate if changes were made.



ABSTRACT

Background: Odontomas are the most common odontogenic tumours and are usually detected during the first two decades of life. They are considered hamartomas rather than true neoplasms and are broadly classified into compound and complex types. Compound odontomas commonly occur in the anterior maxilla and are often associated with delayed eruption or impaction of teeth. In contrast, osteomas are benign osteogenic lesions that differ from odontomas in their histogenesis and clinical behaviour. The occurrence of compound odontomas in the anterior mandible is uncommon.

Case presentation: A 27-year-old female presented with delayed eruption of tooth 33 and displacement of adjacent teeth. Clinical examination revealed a hard swelling in the left anterior mandible. Cone-beam computed tomography (CBCT) revealed multiple radiopaque structures consistent with a compound odontoma. Tooth 32 was found to be impacted near the lesion. Surgical excision of the odontoma and extraction of the affected tooth were performed under local anaesthesia. Histopathological analysis confirmed the diagnosis of a compound odontoma, excluding the possibilities of odontoameloblastoma and ameloblastic fibroodontoma.

Conclusion: This case was notable for the unusual anterior mandibular location of the compound odontoma, deviating from the typical anterior maxillary presentation. There was no history of trauma, infection, or familial predisposition. CBCT played a crucial role in defining the extent of the lesion and its relationship with adjacent structures. Early detection and surgical management facilitated future orthodontic planning for tooth 33. This case highlights the importance of considering compound odontoma in atypical locations when evaluating eruption disturbances. Accurate diagnosis relies on advanced imaging and histopathological confirmation. Prompt intervention prevents developmental complications and supports favourable outcomes.

Keywords: Compound odontome, Cone-beam computed tomography, Enucleation, Mandibular canine impaction, Radiopaque denticles

BACKGROUND

Odontomes are the most frequent odontogenic tumours; however, they are more accurately described as hamartomas because they exhibit restricted growth capacity and nonmalignant behaviour. Initially described by Paul Broca in 1867, the term "odontome" originally encompassed all odontogenic tumours [1,2]. However, with advancements in pathology and diagnostic imaging, this classification has undergone significant modifications. Currently, odontomes are considered to be developmental benign anomalies produced by odontogenic tissue that have both epithelial and ectomesenchymal elements [3,4].

The World Health Organisation's head and neck tumour classification places odontomes in the category of odontogenic tumours involving odontogenic epithelium and odontogenic ectomesenchyme, either with or without hard dental tissues such as enamel and dentin being formed [5]. They are made up of morphologically normal dental tissues but with abnormal structure and organisation that differentiates them from the true neoplasms [6]. Odontomes are classified into two major types based on the morphological appearance: compound and complex odontomes. In the case of complex odontomes, enamel, dentin, and cementum are found but deposited in a disordered mass. In contrast, compound odontomes are made up of a multitude of tooth-like structures that resemble the morphology of normal teeth [7].

Clinically, odontomes are often asymptomatic and are most commonly discovered incidentally during routine radiographic surveys. They can present with signs like delayed eruption of the permanent dentition, retention of primary teeth, or swelling in a localised area. Their development is generally slow and painless, and they seldom cause much discomfort unless infected or encroaching on adjacent structures [4]. Although they are harmless, odontomes can disrupt normal tooth eruption and alignment, particularly in young patients, necessitating early diagnosis and treatment. Various etiologic factors have been postulated in the genesis of odontomes. They include local trauma during dentitional formative years, genetic syndromes including Gardner's syndrome, and odontogenic cell hyperactivity including odontoblasts.

Additionally, gene mutations that interfere with the normal pathway of odontogenesis may be responsible for their formation [5]. The specific pathogenesis remains unclear and is still being investigated. Anatomically, their occurrence is according to their nature. Compound odontomes most often occur in the anterior maxilla, while

complex odontomes usually occur in the posterior mandible [6].

The distribution pattern again assists in their differential diagnosis upon radiographic examinations. Owing to the affinity of compound odontomes for the anterior maxilla, instances within other locations, including the anterior mandible, are regarded as uncommon. This current report presents a rare case of a compound odontome within the anterior mandible of a 27-year-old female, highlighting the need for complete diagnostic examination in rare cases.

CASE PRESENTATION

A 27-year-old female patient reported to the Department of Oral and Maxillofacial Surgery with the chief complaint of tooth pain and mobility in the mandibular anterior region for the last 15 to 20 days. The patient gave a history of occasional dull, diffuse pain and discomfort over the left mandibular anterior region. On clinical examination, the left mandibular canine was found to be missing, and a potential impaction or developmental anomaly was considered. Extraoral examination of the patient did not show facial asymmetry or swelling, nor was lymphadenopathy seen. The intraoral examination revealed grade I mobility in tooth 32 and mild gingival inflammation. To evaluate the cause, an orthopantomogram (OPG) and intraoral periapical radiographs were taken. Radiographic examination demonstrated several small, radiopaque, tooth-shaped structures located in close proximity to the apex of tooth 32. In addition, the impacted left mandibular canine was found at the lower edge of the mandible, indicative of a probable compound odontome. To further assess the lesion and anatomical relationships, a cone-beam computed tomography (CBCT) scan was done. The scan reiterated the existence of multiple calcified tooth-like structures within a well-defined radiolucent cavity.

Notably, the CBCT indicated preservation of bony integrity around the lesion and ascertained that the surrounding neurovascular structures, notably the inferior alveolar nerve, were unaffected. Based on the radiographic and clinical examination, a compound odontome with an impacted left mandibular canine was diagnosed. The odontome was to be surgically enucleated along with the extraction of the affected tooth 32. The operation was performed under local anaesthesia with 2% lignocaine with 1:100,000 epinephrine. Povidone-iodine solution was used to disinfect the perioral region. A rectangular mucoperiosteal flap was raised from the right mandibular lateral incisor to the left first premolar to expose the bone. A bony window was made close to the apex of tooth 72 with a round tungsten carbide bur in a

slow-speed handpiece with continuous irrigation with sterile normal saline to prevent thermal damage. After the removal of the cortical bone, several denticles were seen in the cavity. Seven denticles of differing sizes and shapes, along with the overlying follicular tissue, were removed with caution. Tooth 32, which was pathologically mobile and in close approximation to the lesion, was also removed. The cavity was irrigated well with normal saline to remove any remaining debris. The wound was closed with 3-0 Vicryl sutures after repositioning the mucoperiosteal flap. The post-operative course was smooth, and the patient was instructed on routine post-operative care, including pain management and oral hygiene maintenance. Histopathological examination of the resected tissue was consistent with the diagnosis of compound odontome. The case highlights the significance of early radiographic assessment in diagnosing odontogenic lesions with unusual presentations, as well as the rare occurrence of compound odontomes in the anterior mandible.




Figure 1: Orthopantomogram of the lesion



Figure 2: Incision of the lesion



Figure 3: Exposure of the tooth-like structures



Figure 4: Bony cavity after the mass removal



Figure 5: Specimen after the enucleation

DISCUSSION

Cause of compound odontome development [14]. Still, the origin of this hyperactivity in such cases is yet unknown. One intriguing aspect of odontomas is their significant variety in denticle count. Although earlier reports noted the removal of 4 to 37 denticles, a recent remarkable instance reported that as many as 232 denticles were removed during enucleation [17]. Under local anaesthesia and extraction of tooth 32-related to the odontome, in this case-surgical enucleation was performed. To exclude additional mixed odontogenic neoplasms, such as ameloblastic fibro-odontoma or odontoameloblastoma [18], the sample was sent for histological investigation. Although benign in origin, odontomas can cause disturbances in the regular eruption of teeth. These could manifest as delayed eruption, impaction, over-retention of primary teeth, or malalignment caused by tooth displacement [19]. These consequences underscore the importance of early identification and treatment. The appropriate examination of odontomas has benefited much from radiographic imaging, more especially, from Cone Beam Computed Tomography (CBCT). High-resolution pictures of the lesion and its connection with surrounding anatomical structures in space offered by CBCT maximise diagnosis and surgical planning [20]. For odontomas, surgical excision is the preferred therapy. Small-sized lesions can be eliminated uneventfully, but surgeons should be careful not to damage surrounding structures, particularly in Areas with close anatomical packing. Early intervention ensures superior long-term results by reducing the risk of deformity and promoting the natural eruption of impacted permanent teeth. Regarding the present instance, a follow-up was indicated to evaluate and treat tooth 33, which had not erupted following surgery.

PATIENT PERSPECTIVE

For patients with odontomas, the issue can escalate due to the pain or the mobility of the teeth. One such patient, a 27-year-old female, expressed the discomfort with her left mandibular anterior region. The pain, along with the discomfort, brought about a lot of discomfort and uncertainty concerns to the patient. The radiographic findings of several tooth structures also complicated her case as this would require other forms of imaging like a CT scan to obtain satisfactory clarity to the extent of the issue. Additionally, having a missing tooth and a missing canine caused a lot of self-consciousness and frustration to the patient as well.

When patients were told of having a compound odontoma, and the need to perform an enucleation procedure, it indeed would bring a lot of worries. Even though the procedure was done with the patient under local anesthesia, many of having abnormal growths caused worries such as the recovery procedure. Complications such as the eruption of the impacted 33 tooth would also provide a lot of worries especially with future dental visits. These reasons are understandable, particularly the positive and encompassing emotions. Since there were also worries of a potentially uncomfortable recovery, this might bring the patient to the endwhile worries about the enucleation procedure.

Nevertheless, the anticipated benign results and restoration of regular functionality and the patient's gratifications following the osteosarcoma excision and the scheduled oversight of the affected promontory offer optimism for a constructive subsequent encounter.

CONCLUSION

In essence, odontomas can create major clinical problems if left untreated, even if they are benign and usually asymptomatic. Their varied appearance emphasises the need to include them in differential diagnosis, particularly in younger people with unerupted or impacted teeth. A thorough approach, including clinical, radiological, and histological examinations, ensures appropriate treatment and a favourable prognosis.

REFERENCES

1. Uma E et al. Malays J Med Sci. 2017 May;24(3):92-95. [PMID: 28814937]
2. Patricia GA et al. J Oral Sci. 2011;53(1):67-75. [PMID: 21467817]
3. Sivapathasundharam B et al. J Oral Maxillofac Pathol. 2019 May-Aug;23(2):178-186. [DOI: 10.4103/jomfp.JOMFP_211_19]

4. Boffano P et al. J Craniofac Surg. 2012 May;23(3):685-8. [DOI: 10.1097/SCS.0b013e31824dba1f]
5. Iatrou I et al. J Oral Sci. 2010 Sep;52(3):439-47. [DOI: 10.2334/josnusd.52.439]
6. Philipsen HP et al. Oral Oncol. 1997;33:86-99. [DOI: 10.1016/s0964-1955(96)00067-x]
7. Nelson BL, Thompson LD. Compound odontoma. Head Neck Pathol. 2010;4:290-291. [PMID: 20533004]
8. de Oliveira BH, Campos V, Marçal S. Compound odontoma—diagnosis and treatment: three case reports. Pediatr Dent. 2001;(23):151-157. [PMID: 11340730]
9. Amado Cuesta S et al. Presentation of an erupted complex odontoma. Med Oral. 2003 Nov-Dec;8(5):366-73. [PMID: 14595262].
10. Urvashi S et al. The Saudi Dental Journal. 2010;22:145-149. [PMID: 23960491]
11. Pokharel S, Li Z. Teen gets 232 “teeth” removed in Mumbai. Retrieved January 25, 2025. [Available from: <http://edition.cnn.com/2014/07/25/world/asia/india-abnormal-teeth/>]
12. An S, An C, Choi K. Odontoma: a retrospective study of 73 cases. Imaging Sci Dent. 2012;42:77-81. [PMID: 22783475]
13. Sajesh S et al. J Pharm Bioallied Sci. 2021 Nov;13(Suppl 2):S1772-S1777. [DOI: 10.4103/jpbs.jpbs_270_21]
14. Ahuja VR et al. Int J Clin Pediatr Dent. 2020 Nov-Dec;13(6):729-733. [PMID: 33976503]
15. Marimuthu M et al. Int J Clin Pediatr Dent. 2022 Nov-Dec;15(6):789-792. [PMID: 36866148]
16. Ide F et al. Head Neck Pathol. 2023 Dec;17(4):976-983. [DOI: 10.1007/s12105-023-01593-3]
17. Singh P et al. Ann Maxillofac Surg. 2022 Jan-Jun;12(1):87-90. [DOI: 10.4103/ams.ams_245_21]
18. Chen JW et al. J Am Dent Assoc. 2020 May;151(5):358-367. [DOI: 10.1016/j.adaj.2020.01.015]
19. AlHadidi A et al. Dent Clin North Am. 2024 Apr;68(2):227-245. [PMID: 38417988]
20. Marappa Sathyamoorthy SK et al. BMJ Case Rep. 2021 Sep 20;14(9):e244457. [DOI: 10.1136/bcr-2021-244457]

Ethical Approval: Institutional Review Board approval was not required.

Declaration of Patient Consent: Written informed consent was obtained from the patient prior to initiating the treatment. The patient was fully informed about the nature of the ceramic veneer procedure, the potential risks and benefits, and the expected outcomes. Consent was also obtained for the publication of the case report, including the use of anonymized clinical photographs and information for educational purposes.

Financial Support and Sponsorship: Nil

Conflicts of Interest: The authors declare that they have no conflicts of interest.

Use of Artificial Intelligence (AI) - Assisted Technology for Manuscript Preparation: The authors confirm that no artificial intelligence (AI)- assisted technology was used to assist in the writing or editing of the manuscript, and no images were manipulated using AI tools.

AUTHOR CONTRIBUTIONS:

Vikram Karande-Sanjay Ranade contributed to the conceptualization, writing, data collection, and analysis.

Vaishali Pagare: assisted with study design, data analysis, and manuscript revision.

ABBREVIATIONS USED IN THE STUDY:

- a) **CBCT** – Cone Beam Computed Tomography
- b) **CT** – Computed Tomography
- c) **IARC** – International Agency for Research on Cancer
- d) **WHO** – World Health Organization

DECLARATION ON PUBLICATION ETHICS:

The authors declare that they adhere to the COPE guidelines on publishing ethics, as described elsewhere at <https://publicationethics.org/>. The authors also undertake that they are not associated with any other third party (governmental or non-governmental agencies) linking with any form of unethical issues related to this publication. The authors also declare that they are not withholding any information that is misleading to the publisher regarding this article.

DECLARATION ON OFFICIAL E-MAIL:

The corresponding author declares that a lifetime official e-mail from their institution is not available for all authors.

COMMENTS FROM READERS:

Articles published in the ORAL SPHERE JOURNAL OF DENTAL AND HEALTH SCIENCES are open for relevant post-publication comments and criticisms, which will be published immediately, linking to the original article without open access charges. Comments should be concise, coherent, and critical in fewer than 1000 words.

DISCLAIMER:

The Oral Sphere Journal of Dental and Health Sciences provides a platform for the scholarly communication of data and information to create knowledge in the dental and medical domains after adequate peer/editorial reviews and editing, with entertaining revisions where required. The views and opinions expressed are those of the author(s) and do not reflect the views or opinions of Oral Sphere Journal of Dental and Health Sciences and/or its publisher, Font Fusions Publication Pvt. Ltd. Font Fusions Publication remains neutral and allows authors to specify their address and affiliation details, including territory where required.